Author Index

ABO T. *see* IKARASHI Y.
ABRAMS J. *see* ANDERSSON J.
ACHA-ORBEA H. *see* ANDERSSON M.
ADAMS T.E. *see* WALKER I.D.
ÅGREN K. *see* ANDERSSON J.
AIKAWA M. *see* NAGASAWA H.
AKAZA T. *see* NISHIMURA M.
AKIYAMA S.K. *see* MANDY S.
ALLAN C. *see* RIGAS B.
ALTIN J.G., PAGLER E.B. & PARISH C.R. Evidence for an association of CD45 with 32 000–33 000 MW phosphoproteins on murine T and B lymphocytes, 420
AMBROISE-THOMAS P. *see* PICOT S.
ANDERSSON J., ABRAMS J., BJÖRK L., FUNA K., LITTON M., ÅGREN K. & ANDERSSON U. Concomitant *in vivo* production of 19 different cytokines in human tonsils, 16
ANDERSSON M. & ACHA-ORBEA H. The primary *in vivo* immune response to Mls-1 (*Mtv-7 sag*). Route of injection determines the immune response pattern, 438
ANDERSSON U. *see* ANDERSSON J.
DE ANDRÉS B., CÁRDABA B., DEL POZO V., MARTÍN-OROZCO E., GALLARDO S., TRAMÓN P., PALOMINO P. & LAHOZ C. Modulation of the FcγRII and FcγRIII induced by GM-CSF, IFN-γ and IL-4 on murine eosinophils, 155
APPELBERG R., CASTRO A.G. & SILVA M.T. Neutrophils as effector cells of T-cell-mediated, acquired immunity in murine listeriosis, 302
ARAKAWA M. *see* MITSUMA S.
ARDESTANI S.K. *see* WANG Y.
AVRAMEAS S. *see* DRUET E.
AYALA A. *see* ZHU X.-L.
AYED K. *see* DRUET E.

BAILEY S.W. *see* THORPE S.J.
BANCHEREAU J. *see* FLUCKIGER A.-C.
BATE C.A.W. & KWIATKOWSKI D.P. Stimulators of tumour necrosis factor production released by damaged erythrocytes, 256
BAXTER A.G., HORSFALL A.C., HEALEY D., OZEGBE P., DAY S., WILLIAMS D.G. & COOKE A. Mycobacteria precipitate an SLE-like syndrome in diabetes-prone NOD mice, 227
BECKHAM C. *see* WANG Y.
BENBERNOU N. & NAUCIEL C. Influence of mouse genotype and bacterial virulence in the generation of interferon-γ-producing cells during the early phase of *Salmonella typhimurium* infection, 245
BENJAMIN N. *see* LIVERSIDGE J.
VAN DEN BERG W.B. *see* JACOBS M.J.M.
BERGMEIER L.A., WALKER J., TAO L., CRANAGE M. & LEHNER T. Antibodies to human and non-human primate cellular and culture medium components in macaques vaccinated with the simian immunodeficiency virus, 213
BERNIGAUD A. *see* JOUAULT T.
BIANCO N.E. *see* DE SANCTIS J.B.
BJÖRCK P., BRAESCH-ANDERSEN S. & PAULIE S. Antibodies to distinct epitopes on the CD40 molecule co-operate in stimulation and can be used for the detection of soluble CD40, 430
BJÖRK L. *see* ANDERSSON J.
BLANCA I. *see* DE SANCTIS J.B.
BLANGE I. *see* WAHREN M.
BODMAN K.B. *see* JEDDI P.A.
BORREBAECK C.A.K. *see* MÅRTENSSON C.
BØRRETZEN M. *see* THORPE S.J.
BRAESCH-ANDERSEN S. *see* BJÖRCK P.
BRANDTZAEG P. *see* FARSTAD I.N.
BRASITUS T.A. *see* RIGAS B.
BRATTGJERD S., EVENSEN Ø. & LAUVE A. Effect of injected yeast glucan on the activity of macrophages in Atlantic salmon, *Salmo salar* L., as evaluated by *in vitro* hydrogen peroxide production and phagocytic capacity, 288
BREDIUS R.G.M., FIJEN C.A.P., DE HAAS M., KUIJPER E.J., WEENING R.S., VAN DE WINKEL J.G.J. & OUT T.A. Role of neutrophil FcγRIIa (CD32) and FcγRIIIb (CD16) polymorphic forms in phagocytosis of human IgG1- and IgG3-opsonized bacteria erythrocytes, 624
BREVÉ J. *see* VAN WILSEM E.J.G.
BRIDON J.-M. *see* FLUCKIGER A.-C.
BRIÈRE F. *see* FLUCKIGER A.-C.
BROUWER M. *see* KUIPER H.M.
BURNHAM K. *see* PICKARD S.
BUTTERWORTH A.E. *see* LANGLEY J.G.
BYGBJERG I.C. *see* JAKOBSEN P.H.

CALAME W. *see* DAMOISEAUX J.G.M.C.
CALLOW M.G. *see* HOYNE G.F.
CALVERT J.E. *see* CARSON R.T.
CAMPBELL A.K. *see* TAYLOR K.M.
CANFIELD L.S. *see* MANDY S.
CÁRDABA B. *see* DE ANDRÉS B.
CARLSSON R. *see* MÅRTENSSON C.
CARSON R.T., McDONALD D.F., KEHOE M.A. & CALVERT J.E. Influence of Gm allotype on the IgG subclass response to streptococcal M protein and outer membrane proteins of *Moraxella catarrhalis*, 107
CASADO J.A., MERINO J., CID J., SUBIRA M.L. & SANCHEZ-IBARROLA A. The type of interaction with FcγR in human monocytes determines the efficiency of the generation of oxidative burst, 148
CASTRO A.G. *see* APPELBERG R.
CEREDIG R. *see* TIBERGHIEN F.
CHADEE K. *see* WANG W.
CHAO D. *see* DAMOISEAUX J.G.M.C.
CHAUDRY I.H. *see* ZHU X.-L.
CHÁVEZ R. *see* LASCURAIN P.
CHEN X.-J. & ENERBÄCK L. IgE receptors, IgE content and secretory response of mast cells in athymic rats, 595
CHIANG T.J. *see* HUSSAIN R.
CHOU T. *see* MITSUMA S.
CHUNG S., GORCZYNSKI R., CRUZ B., FINGEROTE R., SKAMENE E., PERLMAN S., LEIBOWITZ J., FUNG L., FLOWERS M. & LEVY G. A Th1 cell line (3E9.1) from resistant A/J mice inhibits induction of macrophage procoagulant activity *in vitro* and protects against MHV-3 mortality *in vivo*, 353
CID J. *see* CASADO J.A.
CLANCY R.L. *see* DUNKLEY M.L.
CLEVERS H.C. *see* WALKER I.D.
COATES J.P. *see* HILL S.
COHEN S.B.A., KATSIKIS P.D., FELDMANN M. & LONDEI M. IL-10 enhances expression of the IL-2 receptor α chain on T cells, 329
COHEN Z. *see* GORCZYNSKI R.M.
ČOLIĆ M., VUČEVIC D., MIYASAKA M., TAMATANI T., PAVLOVIĆ M.D. & DUJIĆ A. Adhesion molecules involved in the binding and subsequent engulfment of thymocytes by a rat thymic epithelial cell line, 449
COOKE A. *see* BAXTER A.G.
CRANAGE M. *see* BERGMEIER L.A.

CRAWFORD S.E. *see* HUANG L.
CRIPPS A.W. *see* DUNKLEY M.L.
CROSSAN J.F. *see* WÓJCIAK B.
CRUZ B. *see* CHUNG S.
CZAJKOWSKA B. *see* MARCINKIEWICZ J.

DAMOISEAUX J.G.M.C., DÖPP E.A., CALAME W., CHAO D., MacPHERSON G.G. & DIJKSTRA C.D. Rat macrophage lysosomal membrane antigen recognized by monoclonal antibody ED1, 140
DANIEL P.B. *see* HOCK B.D.
DASGUPTA J.D. *see* GRANJA C.B.
DAY S. *see* BAXTER A.G.
DEARMAN R.J. *see* HOPE J.C.
DELEMARRE F.G.A., STEVENHAGEN A., KROON F.P., VAN EER M.Y., MEENHORST P.L. & VAN FURTH R. Effect of IFN-γ on the proliferation of *Toxoplasma gondii* in monocytes and monocyte-derived macrophages from AIDS patients, 646
DELVES P.J. *see* JEDDI P.A.
DIJKSTRA C.D. *see* DAMOISEAUX J.G.M.C.
see also KURIMOTO I.
DOCKRELL H.M. *see* HUSSAIN R.
DONADILLE A. *see* PICOT S.
DÖPP E.A. *see* DAMOISEAUX J.G.M.C.
DRUET E., GUERY J.-C., AYED K., GUILBERT B., AVRAMEAS S. & DRUET P. Characteristics of polyreactive and monospecific IgG anti-laminin autoantibodies in the rat mercury model, 489
DRUET P. *see* DRUET E.
DUJIĆ A. *see* CÕLIĆ M.
DUNKLEY M.L., CLANCY R.L. & CRIPPS A.W. A role for $CD4^+$ T cells from orally immunized rats in enhanced clearance of *Pseudomonas aeruginosa* from the lung, 362
DUNNE D.W. *see* LANGLEY J.G.
DWEK R.A. *see* RUDD P.M.

VAN EER M.Y. *see* DELEMARRE F.G.A.
ELHASSAN I.M. *see* JAKOBSEN P.H.
ELSON C.J. *see* GRIFFITHS H.L.
ENERBÄCK L. *see* CHEN X.-J.
ETO M. *see* NISHIMURA Y.
EVENSEN Ø. *see* BRATTGJERD S.

FARSTAD I.N., HALSTENSEN T.S., FAUSA O. & BRANDTZAEG P. Heterogeneity of M-cell-associated B and T cells in human Peyer's patches, 457
FAUSA O. *see* FARSTAD I.N.
FEI K. *see* ROBY K.F.
FELDMANN M. *see* COHEN S.B.A.
FENG Z. *see* MANDY S.
FIJEN C.A.P. *see* BREDIUS R.G.M.
FIJIWARA M. *see* IKARASHI Y.
FINGEROTE R. *see* CHUNG S.
FINKE D. & LIEBERT U.G. $CD4^+$ T cells are essential in overcoming experimental murine measles encephalitis, 184
FLOWERS M. *see* CHUNG S.
FLUCKIGER A.-C., BRIÈRE F., ZURAWSKI G., BRIDON J.-M. & BANCHEREAU J. IL-13 has only a subset of IL-4-like activities on B chronic lymphocytic leukaemia cells, 397
FLYNN J.N. & SILEGHEM M. Involvement of $\gamma\delta$ T cells in immunity to trypanosomiasis, 86
FONTANA A. *see* JUNG S.
FORRESTER J.V. *see* LIVERSIDGE J.
see also MESRI M.
FORTUNE F. *see* RUDD P.M.
FU X.-M. *see* GORCZYNSKI R.M.
FUJIKURA Y. *see* TSUCHIDA M.
FUJIOKA H. *see* NAGASAWA H.
FUKUMOTO T. *see* TSUCHIDA M.
FUNA K. *see* ANDERSSON J.
FUNG L. *see* CHUNG S.
FURRIE E., TURNER M.W. & STROBEL S. Failure of SCID mice to generate an oral tolerogen after a feed of ovalbumin: a role for a functioning gut-associated lymphoid system, 562

VAN FURTH R. *see* DELEMARRE F.G.A.
see also GEERTSMA M.F.
FURUKAWA M. *see* SAKAMOTO A.

GALLAGHER G. *see* WALKER W.
GALLARDO S. *see* DE ANDRÉS B.
GAUS H., MIETHKE T., WAGNER H. & HEEG K. Superantigen-induced anergy of $V\beta8^+$ $CD4^+$ T cells induces functional but non-proliferative T cells *in vivo*, 333
GEERTSMA M.F., ZOMERDIJK T.P.L., NIBBERING P.H. & VAN FURTH R. Pulmonary surfactant inhibits monocyte bactericidal functions by altering activation of protein kinase A and C, 133
GLEW M.D. *see* WALKER I.D.
GODFREY H.P. *see* MANDY S.
GORCZYNSKI R.M., COHEN Z., FU X.-M., LEVY G., PLAPLER H. & SULLIVAN B. Hepatic regulation of lymphocyte adhesion to, and activation on, syngeneic endothelial monolayers, 58
see also CHUNG S.
GOROCICA P. *see* LASCURAIN P.
GOTOH Y. *see* TOKUSHIMA M.
GOZASHTI C.S. *see* GRANJA C.B.
GRABOWSKA A. *see* MARCINKIEWICZ J.
GRABOWSKI P. *see* LIVERSIDGE J.
GRANJA C.B., GOZASHTI C.S. & DASGUPTA J.D. CD4-independent signal transduction through the T-cell receptor (TCR/CD3), 414
GREENWOOD B.M. *see* JAKOBSEN P.H.
GRIFFITHS H.L., KUMPEL B.M., ELSON C.J. & HADLEY A.G. The functional activity of human monocytes passively sensitized with monoclonal anti-D suggests a novel role for FcγRI in the immune destruction of blood cells, 370
GROVES P.L., PECH M.H., TROUTT A.B. & KELSO A. Limiting dilution analysis reveals the precursors of interleukin-4-producing $CD4^+$ cells induced by protein immunization, 25
GRÜWALD C. *see* NEUNER P.
GUERY J.-C. *see* DRUET E.
GUIDA L. *see* HAWRYLOWICZ C.M.
GUILBERT B. *see* DRUET E.

DE HAAS M. *see* BREDIUS R.G.M.
HADLEY A.G. *see* GRIFFITHS H.L.
HALSTENSEN T.S. *see* FARSTAD I.N.
HAMMERSLEY A.P. *see* LANGLEY J.G.
HARA A. *see* MOCHIDA K.
HARDING C.V. & PFEIFER J.D. Antigen expressed by *Salmonella typhimurium* is processed for class I major histocompatibility complex presentation by macrophages but not infected epithelial cells, 670
HARNETT M.M. *see* KIRKHAM P.A.
HART D.N.J. *see* HOCK B.D.
HARTMANN K.-U. *see* ROSENBLATT N.
HARTUNG H.-P. *see* JUNG S.
HAWRYLOWICZ C.M., GUIDA L. & PALEOLOG E. Dexamethasone up-regulates granulocyte–macrophage colony-stimulating factor receptor expression on human monocytes, 274
HEALEY D. *see* BAXTER A.G.
HEATH L.S. *see* JEDDI P.A.
HEEG K. *see* GAUS H.
HEIN W.R. *see* WALKER I.D.
HILL S., COATES J.P., KIMBER I. & KNIGHT S.C. Differential function of dendritic cells isolated from blood and lymph nodes, 295
HIMENO K. *see* NAGASAWA H.
HINSON C.J. *see* MORRISON W.I.
HIROMATSU K. *see* NISHIMURA Y.
HIROSE A. *see* TOKUSHIMA M.
HISAEDA H. *see* NAGASAWA H.
HOCK B.D., STARLING G.C., DANIEL P.B. & HART D.N.J. Characterization of CMRF-44, a novel monoclonal antibody to an activation antigen expressed by the allostimulatory cells within peripheral blood, including dendritic cells, 573
VAN DEN HOEK A.E.M. *see* JACOBS M.J.M.

VAN HOOGSTRATEN I.M.W. *see* VAN WILSEM E.J.G.
HOPE J.C., DEARMAN R.J., KIMBER I. & HOPKINS S.J. The kinetics of cytokine production by draining lymph node cells following primary exposure of mice to chemical allergens, 250
HOPKINS S.J. *see* HOPE J.C.
HORSFALL A.C. *see* BAXTER A.G.
HORUZSKO A., TOMLINSON P.D., STRACHAN T. & MELLOR A.L. Transcription of HLA-G transgenes commences shortly after implantation during embryonic development in mice, 324
HOWARD C.J. *see* MORRISON W.I.
HOYNE G.F., CALLOW M.G., KUO M.-C. & THOMAS W.R. Inhibition of T-cell responses by feeding peptides containing major and cryptic epitopes: studies with the *Der p* I allergen, 190
HSUEH W. *see* HUANG L.
HUANG L., TAN X., CRAWFORD S.E. & HSUEH W. Platelet-activating factor and endotoxin induce tumour necrosis factor gene expression in rat intestine and liver, 65
HUNT J.S. *see* ROBY K.F.
HUSSAIN R., DOCKRELL H.M. & CHIANG T.J. IgG subclass antibody to *Mycobacterium leprae* 18 000 MW antigen is restricted to IgG1 and IgG3 in leprosy, 495
HVIID L. *see* JAKOBSEN P.H.

IIAI T. *see* IKARASHI Y.
IKARASHI Y., ABO T., KAWAI K., IIAI T., WATANABE H., SUZUKI K., MATSUMOTO Y., OMATA S. & FIJIWARA M. Expansion of intermediate T-cell receptor cells in mice with autoimmune-like graft-versus-host disease, 205
ITO K. *see* MITSUMA S.
ITO Y. *see* NAGASAWA H.
IWAMOTO I. *see* SAKAMOTO A.

JACOBS M.J.M., VAN DEN HOEK A.E.M., VAN LENT P.L.E.M., VAN DE LOO F.A.J., VAN DE PUTTE L.B.A. & VAN DEN BERG W.B. Role of IL-2 and IL-4 in exacerbations of murine antigen-induced arthritis, 390
JAKOBSEN P.H., MORRIS-JONES S., RØNN A., HVIID L., THEANDER T.G., ELHASSAN I.M., BYGBJERG I.C. & GREENWOOD B.M. Increased plasma concentrations of sICAM-1, sVCAM-1 and sELAM-1 in patients with *Plasmodium falciparum* or *P. vivax* malaria and association with disease severity, 665
JEDDI P.A., LUND T., BODMAN K.B., SUMAR N., LYDYARD P.M., POUNCEY L., HEATH L.S., KIDD V.J. & DELVES P.J. Reduced galactosyltransferase mRNA levels are associated with the agalactosyl IgG found in arthritis-prone MRL-*lpr*/*lpr* strain mice, 484
JENKINSON E.J. *see* READY A.R.
JOHANSSON P.J.H., OTA T., TSUCHIYA N., MALONE C.C. & WILLIAMS, JR R.C. Studies of protein A and herpes simplex virus-1 induced Fcγ-binding specificities. Different binding patterns for IgG3 from Caucasian and Oriental subjects, 631
DE JONG R. *see* KUIPER H.M.
JONSSON R. *see* WAHREN M.
JOUAULT T., BERNIGAUD A., LEPAGE G., TRINEL P.A. & POULAIN D. The *Candida albicans* phospholipomannan induces *in vitro* production of tumour necrosis factor-α from human and murine macrophages, 268
JUJI T. *see* NISHIMURA M.
JUNG S., SCHLUESENER H.J., SCHMIDT B., FONTANA A., TOYKA K.V. & HARTUNG H.-P. Therapeutic effect of transforming growth factor-β_2 on actively induced EAN but not adoptive transfer EAN, 545

KALLIOMÄKI S. *see* MÅRTENSSON C.
KARIUKI H.C. *see* LANGLEY J.G.
KASINRERK W., MAJDIC O., PRAPUTPITTAYA K. & SITTISOMBUT N. Enhancement of human lymphocyte proliferative response to purified protein derivative by an anti-interleukin-2 receptor α chain antibody (CD25), 33
KASPROWICZ A. *see* MARCINKIEWICZ J.
KATSIKIS P.D. *see* COHEN S.B.A.
KAWAI K. *see* IKARASHI Y.
KEHOE M.A. *see* CARSON R.T.
KELLER K. *see* WANG W.
KELSO A. *see* GROVES P.L.
VAN KESSEL K.P.M. *see* WEERSINK A.J.L.
KHAN M.Y. *see* MANDY S.
KIDD V.J. *see* JEDDI P.A.
KIMBER I. *see* HILL S.
see also HOPE J.C.
KIMOTO M. *see* TOKUSHIMA M.
KIRKHAM P.A., SANTOS-ARGUMEDO L., HARNETT M.M. & PARKHOUSE R.M.E. Murine B-cell activation via CD38 and protein tyrosine phosphorylation, 513
KLOSNER G. *see* NEUNER P.
KNIGHT S.C. *see* HILL S.
KNOBLER R. *see* NEUNER P.
KNOWLES A. *see* SATOH T.
KOARADE S. *see* TOKUSHIMA M.
KOBAYASHI K. *see* OHKURO M.
KOBAYASHI N. *see* NISHIMURA Y.
KOCISZEWSKA B. *see* MARCINKIEWICZ J.
KOIKE T. *see* SAKAMOTO A.
KONG Y.-Y. *see* NISHIMURA Y.
KONISHI M. *see* TSUCHIDA M.
KRAAL G. *see* VAN WILSEM E.J.G.
KRISTENSSON K. *see* MÅRTENSSON C.
KROON F.P. *see* DELEMARRE F.G.A.
KUHRÖBER A., SCHIRMBECK R. & REIMANN J. A self-reactive class I-restricted T-cell response of H-2^b mice to determinants of the Vβ8.2 domain of the T-cell receptor for antigen, 532
KUIJPER E.J. *see* BREDIUS R.G.M.
KUIPER H.M., DE JONG R., BROUWER M., LAMMERS K., WIJDENES J. & VAN LIER R.A.W. Influence of CD28 co-stimulation on cytokine production is mainly regulated via interleukin-2, 38
KUMPEL B.M. *see* GRIFFITHS H.L.
KUNDU S.K. & MERIGAN T.C. Relationship of HIV-1 provirus load, $CD8^+$ $CD11^+$ T cells and HIV-1 envelope-specific cytotoxic T lymphocytes in HIV-infected asymptomatic patients, 81
KUO M.-C. *see* HOYNE G.F.
KURIMOTO I., VAN ROOIJEN N., DIJKSTRA C.D. & STREILEIN J.W. Role of phagocytic macrophages in induction of contact hypersensitivity and tolerance by hapten applied to normal and ultraviolet B-irradiated skin, 281
KWIATKOWSKI D.P. *see* BATE C.A.W.

LAHOZ C. *see* DE ANDRÉS B.
LAMMERS K. *see* KUIPER H.M.
LANGLEY J.G., KARIUKI H.C., HAMMERSLEY A.P., OUMA J.H., BUTTERWORTH A.E. & DUNNE D.W. Human IgG subclass responses and subclass restriction to *Schistosoma mansoni* egg antigens, 651
LASCURAIN P., CHÁVEZ R., GOROCICA P., PÉREZ A., MONTAÑO L.F. & ZENTENO E. Recognition of a $CD4^+$ mouse medullary thymocyte subpopulation by *Amaranthus leucocarpus* lectin, 410
LAUVE A. *see* BRATTGJERD S.
LEHNER T. *see* BERGMEIER L.A.
see also RUDD P.M.
LEIBOWITZ J. *see* CHUNG S.
VAN LENT P.L.E.M. *see* JACOBS M.J.M.
LEPAGE G. *see* JOUAULT T.
LEVY G. *see* CHUNG S.
see also GORCZYNSKI R.M.
LI M.-S. *see* SATOH T.
LIANG B. *see* WANG Y.
LIEBERT U.G. *see* FINKE D.
VAN LIER R.A.W. *see* KUIPER H.M.
LITTON M. *see* ANDERSSON J.
LIVERSIDGE J., GRABOWSKI P., RALSTON S., BENJAMIN N. & FORRESTER J.V. Rat retinal pigment epithelial cells express an inducible form of nitric oxide synthase and produce nitric oxide in response to inflammatory cytokines and activated T cells, 404

see also MESRI M.
LOBO P.I. & PATEL H.C. A novel role for MHC class II antigens: evidence implicating a protective effect on tumour cells against cytotoxicity by NK and LAK cells, 240
LONDEI M. *see* COHEN S.B.A.
VAN DE LOO F.A.J. *see* JACOBS M.J.M.
LOOR F. *see* ROSENBLATT N.
see also TIBERGHIEN F.
LOU Y.-H. *see* MOCHIDA K.
LOWRIE D.B. *see* SILVA C.L.
LUGER T.A. *see* NEUNER P.
LUND T. *see* JEDDI P.A.
LYDYARD P.M. *see* JEDDI P.A.

McDONALD D.F. *see* CARSON R.T.
MACHEINER W. *see* NEUNER P.
MacHUGH N.D. *see* MORRISON W.I.
McKENZIE I.F.C. *see* XING P.-X.
MacPHERSON G.G. *see* DAMOISEAUX J.G.M.C.
MAEDA T. *see* NISHIMURA Y.
MAEKAWA Y. *see* NAGASAWA H.
MAJDIC O. *see* KASINRERK W.
MALONE C.C. *see* JOHANSSON P.J.H.
MANDY K. *see* MANDY S.
MANDY S., FENG Z., CANFIELD L.S., MANDY K., QUAN X., ROWEHL R.A., KHAN M.Y., AKIYAMA S.K. & GODFREY H.P. Inhibition of expression of delayed hypersensitivity by neutralizing monoclonal anti-T-cell fibronectin antibody, 582
MARCINKIEWICZ J., CZAJKOWSKA B., GRABOWSKA A., KASPROWICZ A. & KOCISZEWSKA B. Differential effects of chlorination of bacteria on their capacity to generate NO, TNF-α and IL-6 in macrophages, 611
MÅRTENSSON C., KRISTENSSON K., KALLIOMÄKI S., BORREBAECK C.A.K. & CARLSSON R. Antigen-specific human immunoglobulin production in SCID mice transplanted with human peripheral lymphocytes is dependent on $CD4^+$ $CD45RO^+$ T cells, 171
MARTÍN-OROZCO E. *see* DE ANDRÉS B.
MATSUMOTO Y. *see* IKARASHI Y.
MEENHORST P.L. *see* DELEMARRE F.G.A.
MELLOR A.L. *see* HORUZSKO A.
MERIGAN T.C. *see* KUNDU S.K.
MERINO J. *see* CASADO J.A.
MESRI M., LIVERSIDGE J. & FORRESTER J.V. ICAM-1/LFA-1 interactions in T-lymphocyte activation and adhesion to cells of the blood–retina barrier in the rat, 52
METCALFE S.A. *see* WALKER I.D.
MIETHKE T. *see* GAUS H.
MITOMI Y. *see* NISHIMURA M.
MITSUMA S., YOSHIZAWA H., ITO K., MORIYAMA H., WAKABAYASHI M., CHOU T., ARAKAWA M. & SHU S. Adoptive immunotherapy mediated by anti-TCR/IL-2-activated tumour-draining lymph node cells, 45
MITSUNAGA S. *see* NISHIMURA M.
MIYAKE K. *see* TOKUSHIMA M.
MIYASAKA M. *see* CÕLIĆ M.
MOCHIDA K., LOU Y.-H., HARA A. & YAMAUCHI K. Physical biochemical properties of IgM from a teleost fish, 675
MONTAÑO L.F. *see* LASCURAIN P.
MORGAN A. *see* SMITH R.M.
MORGAN B.P. *see* TAYLOR K.M.
MORIYAMA H. *see* MITSUMA S.
MORRIS-JONES S. *see* JAKOBSEN P.H.
MORRISON M.H. *see* ZHU X.-L.
MORRISON W.I., HOWARD C.J., HINSON C.J., MacHUGH N.D. & SOPP P. Identification of three distinct allelic forms of bovine CD4, 589
MU H.-H. & SEWELL W.A. Regulation of DTH and IgE responses by IL-4 and IFN-γ in immunized mice given pertussis toxin, 639

NAGASAWA H., HISAEDA H., MAEKAWA Y., FUJIOKA H., ITO Y., AIKAWA M. & HIMENO K. γδ T cells play a crucial role in the expression of 65 000 MW heat-shock protein in mice immunized with *Toxoplasma* antigen, 347
NAGASAWA S. *see* OHKURO M.
NAITO K. *see* TSUCHIDA M.
NATVIG J.B. *see* THORPE S.J.
NAUCIEL C. *see* BENBERNOU N.
NEUNER P., KLOSNER G., SCHAUER E., POURMOJIB M., MACHEINER W., GRÜWALD C., KNOBLER R., SCHWARZ A., LUGER T.A. & SCHWARZ T. Pentoxifylline *in vivo* down-regulates the release of IL-1β, IL-6, IL-8 and tumour necrosis factor-α by human peripheral blood mononuclear cells, 262
NIBBERING P.H. *see* GEERTSMA M.F.
NISHIMURA H. *see* TOKUSHIMA M.
NISHIMURA M., MITSUNAGA S., AKAZA T., MITOMI Y., TADOKORO K. & JUJI T. Protection against natural killer cells by interferon-γ treatment of K562 cells cannot be explained by augmented major histocompatibility complex class I expression, 75
NISHIMURA Y., ETO M., MAEDA T., HIROMATSU K., KOBAYASHI N., NOMOTO K., KONG Y.-Y. & NOMOTO K. Inhibition of skin xenograft rejection by depleting T-cell receptor αβ-bearing cells without T-cell receptor γδ-bearing cells or natural killer cells by monoclonal antibody, 196
NOMOTO K. *see* NISHIMURA Y.

O'KEEFFE M.A. *see* WALKER I.D.
OHKURO M., KOBAYASHI K., TAKAHASHI K. & NAGASAWA S. Effect of C1q on the processing of immune complexes by human neutrophils, 507
OLIVEIRA S.C., ZHU Y. & SPLITTER G.A. Recombinant L7/L12 ribosomal protein and γ-irradiated *Brucella abortus* induce a T-helper 1 subset response from murine $CD4^+$ T cells, 659
OMATA S. *see* IKARASHI Y.
OTA T. *see* JOHANSSON P.J.H.
OUMA J.H. *see* LANGLEY J.G.
OUT T.A. *see* BREDIUS R.G.M.
OZEGBE P. *see* BAXTER A.G.

PAGLER E.B. *see* ALTIN J.G.
PALEOLOG E. *see* HAWRYLOWICZ C.M.
PALOMINO P. *see* DE ANDRÉS B.
PAREKH R.B. *see* RUDD P.M.
PARISH C.R. *see* ALTIN J.G.
PARKHOUSE R.M.E. *see* KIRKHAM P.A.
PATEL H.C. *see* LOBO P.I.
PATEL T. *see* RUDD P.M.
PAULIE S. *see* BJÖRCK P.
PAVLOVIĆ M.D. *see* CÕLIĆ M.
PECH M.H. *see* GROVES P.L.
PÉREZ A. *see* LASCURAIN P.
PERLMAN S. *see* CHUNG S.
PETTERSSON I. *see* WAHREN M.
PFEIFER J.D. *see* HARDING C.V.
PICKARD S., SHANKAR G. & BURNHAM K. Langerhans' cell depletion by staphylococcal superantigens, 568
PICOT S., SHEICK I., SYLVI A., DONADILLE A. & AMBROISE-THOMAS P. Signal transduction pathways involved in tumour necrosis factor secretion by *Plasmodium falciparum*-stimulated human monocytes, 70
PIERSON S. *see* PRIOR L.
PIETRO R.C.L.R. *see* SILVA C.L.
PLAPLER H. *see* GORCZYNSKI R.M.
POULAIN D. *see* JOUAULT T.
POUNCEY L. *see* JEDDI P.A.
POURMOJIB M. *see* NEUNER P.
DEL POZO V. *see* DE ANDRÉS B.
PRAPUTPITTAYA K. *see* KASINRERK W.
PRENZOSKA J. *see* XING P.-X.
PRIOR L., PIERSON S. & RIGGS J. Rapid restoration of B-cell function in XID mice by intravenous transfer of peritoneal cavity B cells, 180
VAN DE PUTTE L.B.A. *see* JACOBS M.J.M.

QUAN X. *see* MANDY S.

RADZIOCH D. *see* DE SANCTIS J.B.
RAKHMILEVICH A.L. Evidence that γδ T cells play a limited role in resistance to murine listeriosis, 524
RALSTON S. *see* LIVERSIDGE J.
RANDEN I. *see* THORPE S.J.
READY A.R. & JENKINSON E.J. The role of macrophages and bone marrow-derived dendritic cells in the rejection of fetal thymus allografts, 465
REIMANN J. *see* KUHRÖBER A.
RIGAS B., TSIOULIAS G.J., ALLAN C., WALI R.K. & BRASITUS T.A. The effect of bile acids and piroxicam on MHC antigen expression in rat colonocytes during colon cancer development, 319
RIGGS J. *see* PRIOR L.
ROBY K.F., FEI K., YANG Y. & HUNT J.S. Expression of HLA class II-associated peptide transporter and proteasome genes in human placentas and trophoblast cell lines, 444
VAN ROOIJEN N. *see* KURIMOTO I.
RØNN A. *see* JAKOBSEN P.H.
ROSENBLATT N., HARTMANN K.-U. & LOOR F. The *Yaa* gene-dependent B-cell deficiency worsens the generalized lymphadenopathy and autoimmunity of C57BL/6-*gld* male mice, 476
ROWEHL R.A. *see* MANDY S.
RUDD P.M., FORTUNE F., PATEL T., PAREKH R.B., DWEK R.A. & LEHNER T. A human T-cell receptor recognizes 'O'-linked sugars from the hinge region of human IgA1 and IgD, 99
RUSSELL S. *see* XING P.-X.

SAKAMOTO A., FURUKAWA M., IWAMOTO I., KOIKE T., TOMIOKA H. & SUMIDA T. T-cell receptor Vβ repertoire of L3T4$^+$ regulatory T cells in anti-L3T4 antibody-induced tolerant NOD mice, 540
SANCHEZ-IBARROLA A. *see* CASADO J.A.
DE SANCTIS J.B., BLANCA I., RADZIOCH D. & BIANCO N.E. Lipoprotein lipase expression in natural killer cells and its role in their cytotoxic activity, 232
SANTOS-ARGUMEDO L. *see* KIRKHAM P.A.
SATOH T., KNOWLES A., LI M.-S., SUN L., TOOZE J.A., ZABUCCHI G. & SPRY C.J.F. Expression of lacto-N-fucopentaose III (CD15)- and sialyl-Lewis X-bearing idiopathic hypereosinophilic syndrome, 313
SATOH T., SUN L., LI M.-S. & SPRY C.J.F. Interleukin-5 mRNA levels in blood and bone marrow mononuclear cells from patients with the idiopathic hypereosinophilic syndrome, 308
SCHAUER E. *see* NEUNER P.
SCHEPER R.J. *see* VAN WILSEM E.J.G.
SCHIRMBECK R. *see* KUHRÖBER A.
SCHLUESENER H.J. *see* JUNG S.
SCHMIDT B. *see* JUNG S.
SCHWARZ A. *see* NEUNER P.
SCHWARZ T. *see* NEUNER P.
SEWELL W.A. *see* MU H.-H.
SHANKAR G. *see* PICKARD S.
SHEICK I. *see* PICOT S.
SHIRAI T. *see* TOKUSHIMA M.
SHU S. *see* MITSUMA S.
SILEGHEM M. *see* FLYNN J.N.
SILVA C.L., SILVA M.F., PIETRO R.C.L.R. & LOWRIE D.B. Protection against tuberculosis by passive transfer with T-cell clones recognizing mycobacterial heat-shock protein 65, 341
SILVA M.F. *see* SILVA C.L.
SILVA M.T. *see* APPELBERG R.
SITTISOMBUT N. *see* KASINRERK W.
SKAMENE E. *see* CHUNG S.
SKARSTEIN K. *see* WAHREN M.
SMITH R.M., MORGAN A. & WRAITH D.C. Anti-class II MHC antibodies prevent and treat EAE without APC depletion, 1
SOPP P. *see* MORRISON W.I.
SPLITTER G.A. *see* OLIVEIRA S.C.
SPRY C.J.F. *see* SATOH T.
STARLING G.C. *see* HOCK B.D.
STEVENHAGEN A. *see* DELEMARRE F.G.A.
STRACHAN T. *see* HORUZSKO A.
STREILEIN J.W. *see* KURIMOTO I.
STROBEL S. *see* FURRIE E.
SUBIRA M.L. *see* CASADO J.A.
SULLIVAN B. *see* GORCZYNSKI R.M.
SUMAR N. *see* JEDDI P.A.
SUMIDA T. *see* SAKAMOTO A.
SUN L. *see* SATOH T.
SUZUKI K. *see* IKARASHI Y.
SYLVI A. *see* PICOT S.

TADOKORO K. *see* NISHIMURA M.
TAKAHASHI K. *see* OHKURO M.
TAKAI K. *see* TSUCHIDA M.
TAMATANI T. *see* CÕLIĆ M.
TAN X. *see* HUANG L.
TAO L. *see* BERGMEIER L.A.
TAYLOR K.M., MORGAN B.P. & CAMPBELL A.K. Altered glycosylation and selected mutation in recombinant human complement component C9: effects on haemolytic activity, 501
THEANDER T.G. *see* JAKOBSEN P.H.
THOMAS W.R. *see* HOYNE G.F.
THOMPSON K.M. *see* THORPE S.J.
THORPE S.J., BØRRETZEN M., BAILEY S.W., RANDEN I., NATVIG J.B. & THOMPSON K.M. Human monoclonal rheumatoid factors: incidence of cross-reactions with tissue components and correlation with V_H gene usage, 114
TIBERGHIEN F., CEREDIG R. & LOOR F. Influence of the *lpr* environment on the lymph node cell phenotypes in C57BL/6 *nubg* and *nulpr* chimeras, 552
TOKUSHIMA M., KOARADE S., HIROSE A., GOTOH Y., NISHIMURA H., SHIRAI T., MIYAKE K. & KIMOTO M. *In vivo* induction of IgG anti-DNA antibody by autoreactive mixed haplotype Aβz/Aαd MHC class II molecule-specific CD4$^+$ T-cell clones, 221
VAN DER TOL M.E. *see* WEERSINK A.J.L.
TOMIOKA H. *see* SAKAMOTO A.
TOMLINSON P.D. *see* HORUZSKO A.
TOOZE J.A. *see* SATOH T.
TOYKA K.V. *see* JUNG S.
TRAMÓN P. *see* DE ANDRÉS B.
TRINEL P.A. *see* JOUAULT T.
TROUTT A.B. *see* GROVES P.L.
TSIOULIAS G.J. *see* RIGAS B.
TSUCHIDA M., KONISHI M., TAKAI K., NAITO K., FUJIKURA Y. & FUKUMOTO T. Effects of irradiation, glucocorticoid and FK506 on cell-surface antigen expression by rat thymocytes: a three-colour flow cytofluorometric analysis, 469
TSUCHIYA N. *see* JOHANSSON P.J.H.
TURNER M.W. *see* FURRIE E.

VERHOEF J. *see* WEERSINK A.J.L.
VUĆEVIC D. *see* CÕLIĆ M.

WAGNER H. *see* GAUS H.
WAHREN M., SKARSTEIN K., BLANGE I., PETTERSSON I. & JONSSON R. MRL/*lpr* mice produce anti-Ro 52 000 MW antibodies: detection, analysis of specificity and site of production, 9
WAKABAYASHI M. *see* MITSUMA S.
WALI R.K. *see* RIGAS B.
WALKER I.D., GLEW M.D., O'KEEFFE M.A., METCALFE S.A., CLEVERS H.C., WIJNGAARD P.L.J., ADAMS T.E. & HEIN W.R. A novel multi-gene family of sheep γδ T cells, 517
WALKER J. *see* BERGMEIER L.A.
WALKER W. & GALLAGHER G. The *in vivo* production of specific human antibodies by vaccination of human-PBL-SCID mice, 163
WANG W., KELLER K. & CHADEE K. *Entamoeba histolytica* modulates the nitric oxide synthase gene and nitric oxide production by macrophages for cytotoxicity against amoebae and tumour cells, 601
WANG Y., ARDESTANI S.K., LIANG B., BECKHAM C. & WATSON R.R. Anti-IL-4 monoclonal antibody and IFN-γ administration retards development of immune dysfunction and cytokine dysregulation during murine AIDS, 384

WATANABE H. *see* IKARASHI Y.
WATSON R.R. *see* WANG Y.
WEENING R.S. *see* BREDIUS R.G.M.
WEERSINK A.J.L., VAN KESSEL K.P.M., VAN DER TOL M.E. & VERHOEF J. Modulation of lipopolysaccharide binding to human granulocytes, 617
WIJDENES J. *see* KUIPER H.M.
WIJNGAARD P.L.J. *see* WALKER I.D.
WILLIAMS, JR R.C. *see* JOHANSSON P.J.H.
WILLIAMS D.G. *see* BAXTER A.G.
VAN WILSEM E.J.G., VAN HOOGSTRATEN I.M.W., BREVÉ J., SCHEPER R.J. & KRAAL G. Dendritic cells of the oral mucosa and the induction of oral tolerance. A local affair, 128
VAN DE WINKEL J.G.J. *see* BREDIUS R.G.M.
WÓJCIAK B. & CROSSAN J.F. The effects of T cells and their products on *in vitro* healing of epitenon cell microwounds, 93
WRAITH D.C. *see* SMITH R.M.

XING P.-X., RUSSELL S., PRENZOSKA J. & McKENZIE I.F.C. Discrimination between alternatively spliced STP-A and -B isoforms of CD46, 122

YAMAUCHI K. *see* MOCHIDA K.
YANG Y. *see* ROBY K.F.
YOSHIZAWA H. *see* MITSUMA S.

ZABUCCHI G. *see* SATOH T.
ZELLWEGER R. *see* ZHU X.-L.
ZENTENO E. *see* LASCURAIN P.
ZHU X.-L., AYALA A., ZELLWEGER R., MORRISON M.H. & CHAUDRY I.H. Peritoneal macrophages show increased cytokine gene expression following haemorrhagic shock, 378
ZHU Y. *see* OLIVEIRA S.C.
ZOMERDIJK T.P.L. *see* GEERTSMA M.F.
ZURAWSKI G. *see* FLUCKIGER A.-C.

Subject Index

Adhesion molecules:
  and thymocyte interactions, 449
  in malaria, 665
Agalactosyl IgG:
  association with reduced galactosyltransferase mRNA, 484
AIDS, murine:
  effects of anti-interleukin-4 and interferon-γ administration, 384
Allotype, Gm:
  influence on IgG subclass response, 107
Anti-DNA antibody, IgG:
  induction by $CD4^+$ T-cell clones, 221
Autoantibodies, anti-laminin:
  polyreactive and monospecific, 489

B-cell activation:
  via CD38 and protein tyrosine phosphorylation, 631
Bone marrow-derived cells:
  role in rejection of fetal thymus allografts, 465

C1q:
  effect on neutrophil processing of immune complexes, 507
CD4, bovine:
  identification of three allelic forms, 589
$CD4^+$ $CD45RO^+$ T cells:
  role in human antibody production in SCID mice, 171
$CD4^+$ regulatory T cells in NOD mice, 540
CD15 and sialyl-Lewis X molecules:
  expression and function on human eosinophils, 313
CD25, antibodies to:
  and lymphocyte growth, 33
CD28 co-stimulation:
  influence on cytokine production, 38
CD40:
  stimulation by antibodies to distinct epitopes, 430
CD45:
  association with phosphoproteins, 420
CD46, isoforms:
  discrimination between, 122
Class I major histocompatibility complex presentation of *Salmonella typhimurium*, 670
Complement C9:
  effects of altered glycosylation and selected mutation on haemolytic activity, 501
Cytokine production, *in vivo*:
  in human tonsils, 16
Cytokine production following primary allergen exposure, 250

Delayed hypersensitivity:
  inhibition by anti-T-cell fibronectin antibody, 582
Delayed-type hypersensitivity, and IgE responses to pertussis toxin:
  role of interferon-γ and interleukin-4, 639
Dendritic cells:
  novel activation antigen, 573
Dendritic cells, from blood and lymph nodes:
  differential function, 295

Experimental autoimmunue encephalomyelitis:
  prevention and treatment with anti-major histocompatibility class I antibodies, 1

Fcγ binding specification in Caucasian and Oriental subjects, 513
FcγR:
  functional activity, 370
FcγR allotypes:
  role in polymorphonuclear phagocytosis, 624
FcγRII and III:
  modulation by cytokines, 155

Graft-versus-host disease:
  expansion of intermediate T-cell receptor cells, 205
Granulocyte–macrophage colony-stimulating factor receptor expression, monocytes:
  up-regulation by dexamethasone, 274

HIV infection, virus load:
  $CD8^+$ $CD11^+$ T cells and cytotoxic T lymphocytes in asymptomatic patients, 81
HLA-G transgenes:
  transcription after implantation, 324

IgE receptors and content:
  of mast cells, 595
IgG subclass distribution, anti-*Mycobacterium leprae* 18 000 MW antibodies:
  in leprosy, 495
IgG subclass responses, human:
  to schistosome egg antigen, 651
IgM, from teleost fish:
  physical biochemical properties, 675
Immunotherapy, adoptive:
  mediated by anti-T-cell receptor/interleukin-2 activated T cells, 45
Interferon-γ:
  activation of monocytes and macrophages, 646
  treatment of K562 cells and protection against natural killer cells, 75
Interferon-γ-producing cells:
  influence of mouse genotype and bacterial virulence, 240
Interleukin-2 and interleukin-4:
  role in murine arthritis, 390
Interleukin-4-producing $CD4^+$ cells:
  precursors, 25
Interleukin-5 mRNA:
  in hypereosinophilic syndrome, 308
Interleukin-10:
  enhancement of interleukin-2 receptor α chain on T cells, 329
Interleukin-13:
  interleukin-4-like activities on B cells, 397

Langerhans' cell depletion:
  by superantigen, 568
Lipopolysaccharide binding to human granulocytes, 617
Lipoprotein lipase:
  expression in natural killer cells and role in cytotoxicity, 232
lpr chimeras:
  lymph node phenotypes, 552
Lymphocyte adhesion, to endothelial monolayers:
  hepatic regulation, 58

M-cell-associated B and T cells in human Peyer's patches, 457
Macrophage activity, in Atlantic salmon:
  effect of injected yeast glucan, 288
Macrophage lysosomal membrane antigen, 140
Macrophages:
  cytokine gene expression following shock, 378
  role in hypersensitivity and tolerance, 281
Major histocompatiblity complex antigen expression, during colon cancer development:
  effect of bile acids and piroxicam, 319
Major histocompatibilty complex class II antigens:
  effect on susceptibility of tumour cells to natural killer lysis, 240

Measles encephalitis, experimental murine:
role of $CD4^+$ T cells in overcoming, 184
Mls-1:
in vivo primary response to, 438

Neutrophils:
in cell-mediated immunity against *Listeria*, 302
Nitric oxide:
inhibition by *Entamoeba histolytica*, 601
secretion by macrophages stimulated by chlorinated bacteria, 611

Oral tolerance:
failure in SCID mice, 562
role of dendritic cells, 128
Oxidative burst, in human monocytes:
role of FcγR interaction, 148

Pentoxifylline:
effects on cytokine production, 262
Peptide transporter and proteasome genes:
in trophoblast cells, 444
Phospholipomannan, from *Candida*:
induction of tumour necrosis factor-α, 268
Platelet-aggregation factor:
induction of tumour necrosis factor gene expression, 65
Protein kinase A and C, activation:
alteration by surfactant, 133

Retinal pigment epithelial cells:
expression of nitric oxide synthase, 404
Rheumatoid factors, human, monoclonal:
cross-reactions with tissues and correlation with V_H gene usage, 114
Ro 52 000 MW, antibodies to:
in MRL/*lpr* mice, 9

SCID mice:
production of human antibodies, 163
Self-reactive cytotoxic T lymphocytes, to Vβ8.2 domain of the T-cell receptor, 532
Simian immunodeficiency virus:
antibody to cellular and medium components after immunization of macaques, 213
Superantigen:
Langerhans' cell depletion by, 568
Superantigen-induced energy of $V\beta8^+$ $CD4^+$ T cells, 333
Systemic lupus erythematosus-like syndrome, in NOD mice:
precipitated by mycobacteria, 227

T-cell clones, recognizing heat-shock protein 65:
protection against tuberculosis, 341
T-cell receptor, human:
recognition of 'O'-linked sugars from the hinge of IgA1 and IgD, 99
T-cell receptor signalling:
CD4 independent, 414
T-cell responses:
inhibition by feeding synthetic peptides, 190
T cells:
effect on healing of epitenon cell microwounds, 93
T cells, $CD4^+$:
role in clearance of *Pseudomonas aeruginosa* from lung, 362
γδ T cells:
in immunity to trypanosomiasis, 86
role in expression of 65 000 MW in mice immunized with *Toxoplasma*, 347
role in resistance to listeriosis, 524
γδ T cells, sheep:
multi-gene family, 517
T-lymphocyte activation and adhesion:
ICAM-1–LFA-1 interactions, 52
Th, cell line:
inhibition of procoagulant activity and protection against MHV, 353
Th1 subject response, from murine CD4 T cells:
induced by *Brucella abortus* and ribosomal protein, 659
Thymocyte subpopulation:
recognition by *Amaranthus leucocarpus* lectin, 410
Thymocytes, rat:
cell surface antigen expression, 469
Transforming growth factor-β_2:
effect on experimental autoimmune neuritis, 545
Tumour necrosis factor secretion:
by *Plasmodium falciparum*-stimulated monocytes, 70
Tumour necrosis factor-α, stimulators of:
released by damaged erythrocytes, 256

X-chromosome-linked immunodeficient mice:
B-cell reconstitution, 180
Xenograft rejection, skin:
inhibition by monoclonal antibodies, 196

Yaa gene-dependent B-cell deficiency:
and lymphadenopathy and autoimmunity in *gld* mice, 476